

# Package: ald (via r-universe)

October 31, 2024

**Type** Package

**Title** The Asymmetric Laplace Distribution

**Version** 1.3.1

**Date** 2021-04-04

**Author** Christian E. Galarza <cgalarza88@gmail.com> and Victor H. Lachos <hlachos@ime.unicamp.br>

**Maintainer** Christian E. Galarza <cgalarza88@gmail.com>

**Description** It provides the density, distribution function, quantile function, random number generator, likelihood function, moments and Maximum Likelihood estimators for a given sample, all this for the three parameter Asymmetric Laplace Distribution defined in Koenker and Machado (1999). This is a special case of the skewed family of distributions available in Galarza et.al. (2017) <doi:10.1002/sta4.140> useful for quantile regression.

**License** GPL (>= 2)

**NeedsCompilation** no

**Date/Publication** 2021-04-04 19:10:02 UTC

**Repository** <https://chedgala.r-universe.dev>

**RemoteUrl** <https://github.com/cran/ald>

**RemoteRef** HEAD

**RemoteSha** f97866097faf2f3227110c20fb9451cba64a5e8b

## Contents

ald-package . . . . .	2
ALD . . . . .	3
likALD . . . . .	5
mleALD . . . . .	7
momentsALD . . . . .	8

<b>Index</b>	<b>11</b>
--------------	-----------

---

ald-package

*The Asymmetric Laplace Distribution*

---

## Description

It provides the density, distribution function, quantile function, random number generator, likelihood function, moments and Maximum Likelihood estimators for a given sample, all this for the three parameter Asymmetric Laplace Distribution defined in Koenker and Machado (1999) useful for quantile regression.

## Details

Package: ald  
Type: Package  
Version: 1.0  
Date: 2015-01-27  
License: GPL (>=2)

## Author(s)

Christian E. Galarza <<cgalarza88@gmail.com>> and Victor H. Lachos <<h1lachos@ime.unicamp.br>>

## References

- Koenker, R., Machado, J. (1999). Goodness of fit and related inference processes for quantile regression. *J. Amer. Statist. Assoc.* 94(3):1296-1309.
- Yu, K. & Moyeed, R. (2001). Bayesian quantile regression. *Statistics & Probability Letters*, 54(4), 437-447.
- Yu, K., & Zhang, J. (2005). A three-parameter asymmetric Laplace distribution and its extension. *Communications in Statistics-Theory and Methods*, 34(9-10), 1867-1879.

## See Also

[ALD,momentsALD,likALD,mleALD](#)

## Examples

```
## Let's plot an Asymmetric Laplace Distribution!  
  
##Density  
sseq = seq(-40,80,0.5)  
dens = dALD(y=sseq,mu=50,sigma=3,p=0.75)  
plot(sseq,dens,type="l",lwd=2,col="red",xlab="x",ylab="f(x)", main="ALD Density function")
```

```

## Distribution Function
df = pALD(q=sseq,mu=50,sigma=3,p=0.75)
plot(sseq,df,type="l",lwd=2,col="blue",xlab="x",ylab="F(x)", main="ALD Distribution function")
abline(h=1,lty=2)

##Inverse Distribution Function
prob = seq(0,1,length.out = 1000)
idf = qALD(prob=prob,mu=50,sigma=3,p=0.75)
plot(prob,idf,type="l",lwd=2,col="gray30",xlab="x",ylab=expression(F^{-1}~(x)))
title(main="ALD Inverse Distribution function")
abline(v=c(0,1),lty=2)

#Random Sample Histogram
sample = rALD(n=10000,mu=50,sigma=3,p=0.75)
hist(sample,breaks = 70,freq = FALSE,ylim=c(0,max(dens)),main="")
title(main="Histogram and True density")
lines(sseq,dens,col="red",lwd=2)

## Let's compute the MLE's

param = c(-323,40,0.9)
y = rALD(10000,mu = param[1],sigma = param[2],p = param[3]) #A random sample
res = mleALD(y)

#Comparing
cbind(param,res$par)

#Let's plot

seqq = seq(min(y),max(y),length.out = 1000)
dens = dALD(y=seqq,mu=res$par[1],sigma=res$par[2],p=res$par[3])
hist(y,breaks=50,freq = FALSE,ylim=c(0,max(dens)))
lines(seqq,dens,type="l",lwd=2,col="red",xlab="x",ylab="f(x)", main="ALD Density function")

```

**Description**

Density, distribution function, quantile function and random generation for a Three-Parameter Asymmetric Laplace Distribution defined in Koenker and Machado (1999) useful for quantile regression with location parameter equal to  $\mu$ , scale parameter  $\sigma$  and skewness parameter  $p$ . This is a special case of the skewed family of distributions in Galarza (2016) available in `lqr::SKD`.

**Usage**

```

dALD(y, mu = 0, sigma = 1, p = 0.5)
pALD(q, mu = 0, sigma = 1, p = 0.5, lower.tail = TRUE)
qALD(prob, mu = 0, sigma = 1, p = 0.5, lower.tail = TRUE)
rALD(n, mu = 0, sigma = 1, p = 0.5)

```

**Arguments**

<code>y, q</code>	vector of quantiles.
<code>prob</code>	vector of probabilities.
<code>n</code>	number of observations.
<code>mu</code>	location parameter.
<code>sigma</code>	scale parameter.
<code>p</code>	skewness parameter.
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ otherwise, $P[X > x]$ .

**Details**

If `mu`, `sigma` or `p` are not specified they assume the default values of 0, 1 and 0.5, respectively, belonging to the Symmetric Standard Laplace Distribution denoted by  $ALD(0, 1, 0.5)$ .

As discussed in Koenker and Machado (1999) and Yu and Moyeed (2001) we say that a random variable  $Y$  is distributed as an ALD with location parameter  $\mu$ , scale parameter  $\sigma > 0$  and skewness parameter  $p$  in  $(0,1)$ , if its probability density function (pdf) is given by

$$f(y|\mu, \sigma, p) = \frac{p(1-p)}{\sigma} \exp -\rho_p\left(\frac{y-\mu}{\sigma}\right)$$

where  $\rho_p(\cdot)$  is the so called check (or loss) function defined by

$$\rho_p(u) = u(p - I_{u < 0})$$

, with  $I$  denoting the usual indicator function. This distribution is denoted by  $ALD(\mu, \sigma, p)$  and its  $p$ -th quantile is equal to  $\mu$ .

The scale parameter `sigma` must be positive and non zero. The skew parameter `p` must be between zero and one ( $0 < p < 1$ ).

**Value**

`dALD` gives the density, `pALD` gives the distribution function, `qALD` gives the quantile function, and `rALD` generates a random sample.

The length of the result is determined by `n` for `rALD`, and is the maximum of the lengths of the numerical arguments for the other functions `dALD`, `pALD` and `qALD`.

**Note**

The numerical arguments other than `n` are recycled to the length of the result.

**Author(s)**

Christian E. Galarza <<cgalarza88@gmail.com>> and Victor H. Lachos <<h1achos@ime.unicamp.br>>

## References

- Galarza Morales, C., Lachos Davila, V., Barbosa Cabral, C., and Castro Cepero, L. (2017) Robust quantile regression using a generalized class of skewed distributions. *Stat*,6: 113-130 doi: 10.1002/sta4.140.
- Yu, K., & Zhang, J. (2005). A three-parameter asymmetric Laplace distribution and its extension. *Communications in Statistics-Theory and Methods*, 34(9-10), 1867-1879.

## See Also

[momentsALD](#),[likALD](#),[mleALD](#)

## Examples

```
## Let's plot an Asymmetric Laplace Distribution!

##Density
library(ald)
sseq = seq(-40,80,0.5)
dens = dALD(y=sseq,mu=50,sigma=3,p=0.75)
plot(sseq,dens,type = "l",lwd=2,col="red",xlab="x",ylab="f(x)", main="ALD Density function")

#Look that is a special case of the skewed family in Galarza (2017)
# available in lqr package, dSKD(...,sigma = 2*3,dist = "laplace")

## Distribution Function
df = pALD(q=sseq,mu=50,sigma=3,p=0.75)
plot(sseq,df,type="l",lwd=2,col="blue",xlab="x",ylab="F(x)", main="ALD Distribution function")
abline(h=1,lty=2)

##Inverse Distribution Function
prob = seq(0,1,length.out = 1000)
idf = qALD(prob=prob,mu=50,sigma=3,p=0.75)
plot(prob,idf,type="l",lwd=2,col="gray30",xlab="x",ylab=expression(F^{-1}(x)))
title(main="ALD Inverse Distribution function")
abline(v=c(0,1),lty=2)

#Random Sample Histogram
sample = rALD(n=10000,mu=50,sigma=3,p=0.75)
hist(sample,breaks = 70,freq = FALSE,ylim=c(0,max(dens)),main="")
title(main="Histogram and True density")
lines(sseq,dens,col="red",lwd=2)
```

---

likALD

*Log-Likelihood function for the Asymmetric Laplace Distribution*

---

## Description

Log-Likelihood function for the Three-Parameter Asymmetric Laplace Distribution defined in Koenker and Machado (1999) useful for quantile regression with location parameter equal to  $\mu$ , scale parameter  $\sigma$  and skewness parameter  $p$ .

**Usage**

```
likALD(y, mu = 0, sigma = 1, p = 0.5, loglik = TRUE)
```

**Arguments**

y	observation vector.
mu	location parameter $\mu$ .
sigma	scale parameter $\sigma$ .
p	skewness parameter $p$ .
loglik	logical; if TRUE (default), the Log-likelihood is return, if not just the Likelihood.

**Details**

If mu, sigma or p are not specified they assume the default values of 0, 1 and 0.5, respectively, belonging to the Symmetric Standard Laplace Distribution denoted by  $ALD(0, 1, 0.5)$ .

As discussed in Koenker and Machado (1999) and Yu and Moyeed (2001) we say that a random variable Y is distributed as an ALD with location parameter  $\mu$ , scale parameter  $\sigma > 0$  and skewness parameter  $p$  in  $(0,1)$ , if its probability density function (pdf) is given by

$$f(y|\mu, \sigma, p) = \frac{p(1-p)}{\sigma} \exp -\rho_p\left(\frac{y-\mu}{\sigma}\right)$$

where  $\rho_p(\cdot)$  is the so called check (or loss) function defined by

$$\rho_p(u) = u(p - I_{u < 0})$$

, with  $I$ , denoting the usual indicator function. Then the Log-likelihood function is given by

$$\sum_{i=1}^n \log\left(\frac{p(1-p)}{\sigma} \exp -\rho_p\left(\frac{y_i - \mu}{\sigma}\right)\right)$$

The scale parameter sigma must be positive and non zero. The skew parameter p must be between zero and one ( $0 < p < 1$ ).

**Value**

likeALD returns the Log-likelihood by default and just the Likelihood if loglik = FALSE.

**Author(s)**

Christian E. Galarza <<cgalarza88@gmail.com>> and Victor H. Lachos <<h1lachos@ime.unicamp.br>>

## References

- Koenker, R., Machado, J. (1999). Goodness of fit and related inference processes for quantile regression. *J. Amer. Statist. Assoc.* 94(3):1296-1309.
- Yu, K. & Moyeed, R. (2001). Bayesian quantile regression. *Statistics & Probability Letters*, 54(4), 437-447.
- Yu, K., & Zhang, J. (2005). A three-parameter asymmetric Laplace distribution and its extension. *Communications in Statistics-Theory and Methods*, 34(9-10), 1867-1879.

## See Also

[ALD](#), [momentsALD](#), [mleALD](#)

## Examples

```
## Let's compute the log-likelihood for a given sample

y = rALD(n=1000)
loglik = likALD(y)

#Changing the true parameters the loglik must decrease
loglik2 = likALD(y,mu=10,sigma=2,p=0.3)

loglik;loglik2
if(loglik>loglik2){print("First parameters are Better")}
```

---

mleALD	<i>Maximum Likelihood Estimators (MLE) for the Asymmetric Laplace Distribution</i>
--------	--

---

## Description

Maximum Likelihood Estimators (MLE) for the Three-Parameter Asymmetric Laplace Distribution defined in Koenker and Machado (1999) useful for quantile regression with location parameter equal to  $\mu$ , scale parameter  $\sigma$  and skewness parameter  $p$ .

## Usage

```
mleALD(y, initial = NA)
```

## Arguments

<code>y</code>	observation vector.
<code>initial</code>	optional vector of initial values $c(\mu, \sigma, p)$ .

## Details

The algorithm computes iteratively the MLE's via the combination of the MLE expressions for  $\mu$  and  $\sigma$ , and then maximizing with respect to  $p$  the Log-likelihood function (`likALD`) using the well known `optimize` R function. By default the tolerance is  $10^{-5}$  for all parameters.

**Value**

The function returns a list with two objects

`iter`                iterations to reach convergence.  
`par`                 vector of Maximum Likelihood Estimators.

**Author(s)**

Christian E. Galarza <<cgalarza88@gmail.com>> and Victor H. Lachos <<hlachos@ime.unicamp.br>>

**References**

Yu, K., & Zhang, J. (2005). A three-parameter asymmetric Laplace distribution and its extension. *Communications in Statistics-Theory and Methods*, 34(9-10), 1867-1879.

**See Also**

[ALD](#), [momentsALD](#), [likALD](#)

**Examples**

```
## Let's try this function

param = c(-323,40,0.9)
y = rALD(10000,mu = param[1],sigma = param[2],p = param[3]) #A random sample
res = mleALD(y)

#Comparing
cbind(param,res$par)

#Let's plot

seqq = seq(min(y),max(y),length.out = 1000)
dens = dALD(y=seqq,mu=res$par[1],sigma=res$par[2],p=res$par[3])
hist(y,breaks=50,freq = FALSE,ylim=c(0,max(dens)))
lines(seqq,dens,type="l",lwd=2,col="red",xlab="x",ylab="f(x)", main="ALD Density function")
```

**Description**

Mean, variance, skewness, kurtosis, central moments w.r.t mu and first absolute central moment for the Three-Parameter Asymmetric Laplace Distribution defined in Koenker and Machado (1999) useful for quantile regression with location parameter equal to mu, scale parameter sigma and skewness parameter p.



**Usage**

```

meanALD(mu=0, sigma=1, p=0.5)
varALD(mu=0, sigma=1, p=0.5)
skewALD(mu=0, sigma=1, p=0.5)
kurtALD(mu=0, sigma=1, p=0.5)
momentALD(k=1, mu=0, sigma=1, p=0.5)
absALD(sigma=1, p=0.5)

```

**Arguments**

k	moment number.
mu	location parameter $\mu$ .
sigma	scale parameter $\sigma$ .
p	skewness parameter $p$ .

**Details**

If mu, sigma or p are not specified they assume the default values of 0, 1 and 0.5, respectively, belonging to the Symmetric Standard Laplace Distribution denoted by  $ALD(0, 1, 0.5)$ .

As discussed in Koenker and Machado (1999) and Yu and Moyeed (2001) we say that a random variable Y is distributed as an ALD with location parameter  $\mu$ , scale parameter  $\sigma > 0$  and skewness parameter  $p$  in  $(0,1)$ , if its probability density function (pdf) is given by

$$f(y|\mu, \sigma, p) = \frac{p(1-p)}{\sigma} \exp -\rho_p\left(\frac{y-\mu}{\sigma}\right)$$

where  $\rho_p(\cdot)$  is the so called check (or loss) function defined by

$$\rho_p(u) = u(p - I_{u < 0})$$

, with  $I$  denoting the usual indicator function. This distribution is denoted by  $ALD(\mu, \sigma, p)$  and its  $p$ th quantile is equal to  $\mu$ . The scale parameter sigma must be positive and non zero. The skew parameter p must be between zero and one ( $0 < p < 1$ ).

**Value**

meanALD gives the mean, varALD gives the variance, skewALD gives the skewness, kurtALD gives the kurtosis, momentALD gives the  $k$ th central moment, i.e.,  $E(y - \mu)^k$  and absALD gives the first absolute central moment denoted by  $E|y - \mu|$ .

**Author(s)**

Christian E. Galarza <<cgalarza88@gmail.com>> and Victor H. Lachos <<h1lachos@ime.unicamp.br>>

**References**

- Koenker, R., Machado, J. (1999). Goodness of fit and related inference processes for quantile regression. *J. Amer. Statist. Assoc.* 94(3):1296-1309.
- Yu, K. & Moyeed, R. (2001). Bayesian quantile regression. *Statistics & Probability Letters*, 54(4), 437-447.
- Yu, K., & Zhang, J. (2005). A three-parameter asymmetric Laplace distribution and its extension. *Communications in Statistics-Theory and Methods*, 34(9-10), 1867-1879.

**See Also**

[ALD,likALD,mleALD](#)

**Examples**

```
## Let's compute some moments for a Symmetric Standard Laplace Distribution.

#Third raw moment
momentALD(k=3,mu=0,sigma=1,p=0.5)

#The well known mean, variance, skewness and kurtosis
meanALD(mu=0,sigma=1,p=0.5)
varALD(mu=0,sigma=1,p=0.5)
skewALD(mu=0,sigma=1,p=0.5)
kurtALD(mu=0,sigma=1,p=0.5)

# and this guy
absALD(sigma=1,p=0.5)
```

# Index

- \* **ALD**
    - ALD, 3
    - likALD, 5
    - mleALD, 7
    - momentsALD, 8
  - \* **Laplace**
    - ALD, 3
    - likALD, 5
    - mleALD, 7
    - momentsALD, 8
  - \* **Log-likelihood**
    - likALD, 5
  - \* **MLE**
    - mleALD, 7
  - \* **Maximum likelihood estimators**
    - mleALD, 7
  - \* **asymmetric laplace distribution**
    - ALD, 3
    - likALD, 5
    - mleALD, 7
    - momentsALD, 8
  - \* **likelihood**
    - likALD, 5
  - \* **moments**
    - momentsALD, 8
  - \* **package**
    - ald-package, 2
  - \* **quantile regression**
    - ALD, 3
    - likALD, 5
    - mleALD, 7
    - momentsALD, 8
- absALD (momentsALD), 8  
ALD, 2, 3, 7, 8, 10  
ald (ald-package), 2  
ald-package, 2
- dALD (ALD), 3
- kurtALD (momentsALD), 8
- likALD, 2, 5, 5, 8, 10
- meanALD (momentsALD), 8  
mleALD, 2, 5, 7, 7, 10  
momentALD (momentsALD), 8  
momentsALD, 2, 5, 7, 8, 8
- pALD (ALD), 3
- qALD (ALD), 3
- rALD (ALD), 3
- skewALD (momentsALD), 8
- varALD (momentsALD), 8